


Preliminary findings concerning emergence and time of beginning and completion of stages of the introduced southern potato wireworm in California compare favorably to findings in the southeastern states. The species appears to be well adapted climatically to this area and only time will determine its destructiveness.

ACKNOWLEDGMENTS

My thanks to J. Wilcox for his interest and helpful suggestions and for operating a light trap and collecting elaterids at Olive (Orange Co.). Thanks also to Saul Frommer, University of California, Riverside, for his critical review of the manuscript.

LITERATURE CITED

- DAY, A., F. P. CUTHBERT, JR., AND W. J. REID. 1971. The southern potato wireworm. USDA Tech. Bull. 1443:1-33.
- REID, W. J., JR. AND CUTHBERT, F. P., JR. 1956. Resistance of the southern potato wireworm to insecticides. J. Econ. Ent. 49:879-880.
- STONE, M. W. 1975. Distribution of four introduced *Conoderus* species in California. *Elateridae*. Coleop. Bull. 29(3):163.



COLEOPTERIST NEWSLETTER

The **Newsletter** has been reinstated by a very large majority of the voting membership (134 in favor, 13 against).

Once again I ask for your support in making the **Newsletter** the kind you want. Send articles and news to be included, eg. techniques, field trip notes, general notes on biology, news about Coleopterists, etc.

We will publish in the *Bulletin*, with the **Newsletter** clearly labelled. Frequency will depend on amount of material submitted. I shall look forward to hearing from you as it is your **Newsletter** and depends entirely on you and your support.

—Charles W. O'Brien, Editor
Laboratory of Aquatic Entomology,
P. O. Box 111,
Florida A and M University,
Tallahassee, Florida 32307

